arXiv:1210.4451v1 [math.CO] 16 Oct 2012

LINEAR EMBEDDINGS OF GRAPHS AND GRAPH LIMITS

HUDA CHUANGPISHIT, MAHYA GHANDEHARI, MATT HURSHMAN,
JEANNETTE JANSSEN, AND NAUZER KALYANIWALLA

Abstract. Many real-life networks can be modelled by stochastic processes with a spatial embedding. In such processes, the link probability decreases with distance. Using the theory of graph limits, we show how to recognize graph sequences produced by random graphs with a linear embedding (a natural embedding into $\mathbb{R}$). We define an operator Γ which applies to graph limits, which assumes a value zero precisely for graph limits of random graphs with a linear embedding. Moreover we introduce a corresponding graph parameter Γ^* and show that, for graph sequences which converge to a graph limit under the cut-norm, the Γ^*-values converge to the Γ-value of the limit. Thus, graph sequences that have diminishing Γ^*-values can be seen as the result of a random graph process with a linear embedding. Finally, we investigate random graphs with a uniform linear embedding, where the probability of a link between two points only depends on their distance.

1. Introduction

The principal purpose of modelling real-life networks is to be able to extract information about these networks. If one assumes such networks are the manifestation of an underlying reality, then a useful way to model these networks is to take a latent space approach. In this approach, the formation of the graph is informed by the hidden spatial reality. The graph formation is modelled as a stochastic process, where the probability of a link occurring between two vertices decreases as their metric distance increases.

The spatial reality can be used to represent attributes of the vertices which are inaccessible or unknown, but which are assumed to inform link formation. For example, in a social network, vertices may be considered as members of a social space, where the coordinates represent the interests and background of the users. Given only the graph, such a spatial model allows us to mine the underlying spatial reality. This approach was taken by Hoff *et al.* in [18]. Recent spatial models for web-based networks, where the vertices clearly exist in an underlying space can be found in [5, 10, 19], while protein-protein interaction networks are modelled as geometric graphs in [23]. A one-dimensional spatial model, the *niche model*, is proposed in [26] to model food webs.

Date: October 17, 2012.

1991 *Mathematics Subject Classification.* Primary 46L07, 47B47.

Key words and phrases. Graph limits, geometric graphs, spatial graph models, linear embedding, random graphs.

The authors wish to thank AARMS, MPrime and NSERC for supporting this work. Ghandehari and Janssen acknowledge the American Institute of Mathematics (AIM) for an invitation to attend the workshop on graph and hypergraph limits, August 2011.

A fundamental question is to determine whether a given network structure, obtained from a real-life network, is compatible with a spatial model. That is, given a graph, how can we judge whether this graph is likely generated by a spatial model, and if so, what are the dimension and other characteristics of the underlying metric space? This question can be answered if a sequence of graphs (snap shots of the stochastic process) is available.

Past efforts have used graph sequences to model dynamic data (see for example [13, 20]). More recently, graph sequences themselves have become a field of active study, where the notion of convergence and the nature of the limit of such graph sequences are developed. A theory of graph limits for sequences of dense graphs has emerged.

In the theory of dense graph limits, put forth by Lovász and Szegedy in [21], convergence is based on homomorphism densities, and the limit is a symmetric function. The theory was developed and extended to sequences of random graphs by Borgs *et al.* in [6, 7, 9] and is being explored further by Lovász and co-authors, and others (see for example [8, 11, 22]). As shown by Diaconis and Janson in [15], the theory of graph limits is closely connected to the probabilistic theory of exchangeable arrays. A different view, where the limit object is referred to as a *kernel*, is provided by Bollobás, Janson and Riordan in [1, 2]. The connection with the results of Borgs *et al.* and an extension of the theory to sparse graphs are presented in [4].

The approach we take in this paper to characterize graphs with a spatial embedding is based on a paper by Bollobás, Janson and Riordan on monotone graph limits. In this paper, a graph parameter Ω is introduced which assumes value zero only for threshold graphs. It is then shown that sequences of graphs for which Ω converges to zero have a limit that is a *monotone* function. Thus, monotone graph limits can be seen as generalizations of threshold graphs.

Diaconis, Holmes and Janson also consider the limits of threshold graphs, see [15]. In [25], they similarly consider the limits of interval graphs. Note that the one-dimensional geometric graphs mentioned in our paper are a special class of interval graphs, namely unit interval or proper interval graphs. However, the authors of [25] focus on different properties and generalizations of interval graphs, and their results do not apply to the problems we consider here.

We show that examining the limit of a graph sequence allows us to conclude, with high probability, if the graphs have a spatial embedding. In this work we only consider random graphs with linear embeddings *i.e.* embeddings in $\mathbb{R}$. We consider embeddings in $\mathbb{R}$ in the most general sense: A graph has a linear embedding if, for a vertex v, and a constant $a \in [0,1]$, all the points with link probability to v that is greater than a lie in an interval around v. We also consider the case of uniform embeddings, where the link probability to a vertex only depends upon the distance from the vertex.

Given a sequence of graphs $\{G_n\}$ with graph limit w, we introduce techniques to use w to recognize whether the graphsG_n have a linear embedding. This is explained in Section 3. We introduce a new graph parameter Γ^*, and its continuous analogue, Γ, which applies to symmetric measurable functions. We show that the limit of the sequence $\{\Gamma^*(G_n)\}$ is closely related to $\Gamma(w)$. In particular $\Gamma^*(G_n) \to 0$ as $n \to \infty$ if and only if w has a linear embedding. In Section 2, we briefly review the results from the theory of graph limits. Finally, Section 5 contains proofs too long to be included in the main body of the paper.

2. Preliminaries: graph limits

In this section we summarize the basic definitions and results from the theory of graph limits, insofar as they are relevant to this paper. For more background, the reader is referred to the papers referenced in the introduction. In this section, we follow the terminology of [21].

Let F and G be two simple graphs, let $hom(F,G)$ be the number of homomorphisms of F into G. The homomorphism density of F into G is defined as

$$t(F,G) = \frac{hom(F,G)}{|V(G)|^{|V(F)|}}.$$

The homomorphism density can be interpreted as the probability that a random mapping $V(F) \to V(G)$ is a homomorphism.

Let G_n be a sequence of simple graphs such that $|V(G_n)| \to \infty$. We can define a notion of convergence based on homomorpism densities.

Definition 2.1. *We say that the sequence* $\{G_n\}$ *converges if for every simple graph* F*, the sequence* $\{t(F,G_n)\}$ *converges.*

This definition of convergence is non-trivial only for dense graphs, *i.e.* for graph sequences $\{G_n\}$ with the property that $|E(G_n)| = \Omega(|V(G_n)|^2)$. When $\{G_n\}$ consists of sparse graphs, then for all graphs F with at least one edge, $t(F,G_n) \to 0$.

A natural metric space to define the convergence of a graph sequence and its limit object is the space of all symmetric, measurable functions from $[0,1]^2$ to $[0,1]$. We will denote this space by $\mathcal{W}_0$. We also denote by $\mathcal{W}$ the space of all the bounded symmetric measurable functions from $[0,1]^2$ to $\mathbb{R}$. We can extend the definition of homomorphism densities to $\mathcal{W}$ as follows. For each function $w \in \mathcal{W}$, let

$$t(F,w) = \int_{[0,1]^k} \prod_{ij \in E(F)} w(x_i,x_j)dx_1 \dots dx_k \tag{1}$$

where $V(F) = \{1,2,\dots,k\}$.

A simple, finite graph G, with adjacency matrix A and with $V(G) = \{1,2,\dots,n\}$, can be represented by a function $w_G \in \mathcal{W}_0$, which takes values in $\{0,1\}$. Split the interval $[0,1]$ into n equal intervals $I_1, I_2 \dots, I_n$. Now for $(x,y) \in I_i \times I_j$,

$$w_G(x,y) = \begin{cases} A_{i,j} & \text{for } i \neq j \\ 1, & \text{for } i = j \end{cases} \tag{2}$$

Our definition of w_G differs slightly from that given in [21] since we give the diagonal blocks $I_i \times I_i$ value one, not zero. This choice was made to preserve the "diagonally increasing" nature of the function. It is a convenience and is not essential for the results.

Note that a graph could be represented by many different functions w_G. Each labelling of the vertices of G, would result in a permutation of the rows and columns of the adjacency matrix, and lead to a trivially different function. In fact, since a graph represents an entire isomorphism class, we need to introduce an equivalent notion for functions in $\mathcal{W}$. Let Φ be the set of all invertible maps $\phi : [0,1] \to$

$[0,1]$ such that both ϕ and is inverse are measure preserving. Any $\phi \in \Phi$ acts on a function $w \in \mathcal{W}$ by transforming it into a function w^{ϕ}, where $w^{\phi}(x,y) = w(\phi(x), \phi(y))$.

The notion of the convergence of a graph sequence can be better understood if $\mathcal{W}$ is equipped with a metric derived from the cut-norm, introduced in [16] and defined as follows. For all $w \in \mathcal{W}$,

$$\|w\|_{\square} = \sup_{S,T \subset [0,1]} \Big| \int_{S \times T} w(x,y)\,dx\,dy \Big|. \tag{3}$$

The cut distance is a metric on the quotient space $\mathcal{W}/\Phi$ derived from the cut norm as follows:

$$\delta_{\square}(w_1, w_2) = \inf_{\phi \in \Phi} \|w_1 - w_2^{\phi}\|_{\square} = \inf_{\phi \in \Phi} \sup_{S,T \subset [0,1]} \Big| \int_{S \times T} (w_1 - w_2^{\phi}) \Big|. \tag{4}$$

It was shown in Theorem 2.3 in [9] that a graph sequence $\{G_n\}$ converges whenever the corresponding sequence of functions w_{G_n} is $\delta_{\square}$-Cauchy. Therefore, the limit is represented by a function in $\mathcal{W}_0$ (not necessarily integer-valued, or corresponding to a graph). The homomorphism densities $t(F, G_n)$ converge to the homomorphism densities $t(F, w)$, and thus the function w (and the isomorphism class it represents) encodes the common structure of the graphs of the sequence. For more details, see [21].

Finally, given any function $w \in \mathcal{W}_0$, there exists a random graph sequence $G(n, w)$ which almost surely converges to w. The random graph $G(n, w)$ has vertex set $\{1, 2, \ldots, n\}$. Each vertex j receives a value x_j, drawn independently and uniformly at random from $[0,1]$. For each pair $i < j$, independently, the probability that vertices i and j are linked equals $w(x_i, x_j)$.

3. Linear embeddings and the parameter Γ^*

In this section, we show how to recognize graph sequences produced by a random graph process with a linear embedding. To this end, we will produce a graph parameter Γ^*, and a related parameter Γ which applies to functions in $\mathcal{W}_0$. Using graph limits, we will show a close relationship between linear embeddings and graph sequences for which Γ^* goes to zero.

First, we need precise definitions of the concepts discussed in the introduction. While this paper is about graphs that can be embedded in the one-dimensional space $(\mathbb{R}, |\cdot|)$, for completeness we extend our definition to general metric spaces. The notation $u \sim v$ signifies "u is adjacent to v".

Definition 3.1. *A random graph on a vertex set V has a spatial embedding into a metric space (S, d) if and only if there exists an injection $\pi : V \to S$ such that for every $a \in [0,1]$ and for every $v \in V$, there exists a convex region $R_a(v)$ of S such that $\pi(v) \in R_a(v)$, and for all vertices $u \in V - \{v\}$, $\pi(u) \in R_a(v)$ if and only if $Prob(u \sim v) > a$. A linear embedding is a spatial embedding into $(\mathbb{R}, |\cdot|)$.*

The notion of spatial embedding can be seen as a "randomized" version of a geometric graph. A graph G is called a *geometric graph* on a metric space (S, d) if there exists an embedding π of the vertices of G in S such that for every two vertices u and v of G, u is adjacent to v if and only if $d(u, v) \leq 1$. Geometric graphs have been studied extensively; see for example [12, 14, 24]. In this paper, we restrict ourselves to geometric graphs on the one dimensional space $(\mathbb{R}, |\cdot|)$ and will refer to these as one-dimensional geometric graphs. If we interpret a graph as a random graph with link probabilities equal to zero or one, then clearly a geometric graph has a spatial embedding.

We introduce first a graph parameter Γ^*, which characterizes geometric graphs in $(\mathbb{R}, |\cdot|)$. One-dimensional geometric graphs are also known as unit interval graphs. The correspondence becomes clear if we associate each vertex u of a one-dimensional geometric graph with the interval $[\pi(u) - \frac{1}{2}, \pi(u) + \frac{1}{2}]$, where π is the geometric embedding. Now vertices u and v are adjacent precisely when the associated intervals overlap.

It is well known that unit interval graphs are characterized by the consecutive 1s property of the vertex-clique matrix (see [17]). Restating this property, it follows that a graph G is one-dimensional geometric if and only if there exists an ordering $\prec$ on the vertex set of G such that

$$\forall v, z, w \in V(G),\ v \prec z \prec w \ \text{ and } \ v \sim w \Rightarrow \ z \sim v \ \text{ and } \ z \sim w. \tag{5}$$

To be self-contained, we present a direct proof below.

Proposition 3.2. *A graph G is a one-dimensional geometric graph if and only if there exists an ordering $\prec$ on $V(G)$ that satisfies (5).*

Proof. The "if" direction is clear. To prove the converse, we proceed by induction. Suppose that for every graph G with $k < n$ vertices, if $V(G)$ satisfies (5) for an ordering $\prec$, then there exists a linear embedding π of vertices of G, with the additional conditions that π is injective, and that the distance between adjacent vertices is strictly less than one. Also, we assume that the embedding respects the ordering $\prec$, so $u \prec v$ implies that $\pi(u) < \pi(v)$.

Suppose that G is a graph with n vertices, and there exists an ordering $\prec$ on vertices of G which satisfies (5).

Let $\{v_1, \ldots, v_n\}$ be the vertices of G labeled such that $v_i \prec v_j$ whenever $i < j$. The ordering $\prec$ restricted to $V(G) \setminus \{v_n\}$ satisfies Condition (5) for $G - v_n$. Thus, by the induction hypothesis, $G - v_n$ has a linear embedding π of $V(G) \setminus \{v_n\}$ into the real line which satisfies the additional conditions. Suppose that m is the smallest index such that v_m is adjacent to v_n. Let $\ell = \max\{\pi(v_{n-1}), \pi(v_{m-1}) + 1\}$, and consider the interval $(\ell, \pi(v_m) + 1)$. By the induction hypothesis, $\pi(v_{m-1}) < \pi(v_m)$, and, since v_m and v_n are adjacent, so are v_m and v_{n-1}, and thus $\pi(v_{n-1}) < \pi(v_m) + 1$. This implies that $\ell < \pi(v_m) + 1$, and thus the interval is non-empty. Moreover, every point in the interval has distance greater than one to all embeddings of non-neighbours of v_n, and distance less than one to all embeddings of neighbours of v_n. Therefore, choosing $\pi(v_n)$ in this interval results in a linear embedding of $V(G)$ with the desired properties, and we are done. $\square$

Using Condition (5), we define a parameter Γ^* on graphs which identifies the one-dimensional geometric graphs. Let G be a graph with a linear order $\prec$ on its vertices. For every $v \in V(G)$, we define the down-set $D(v)$ and the up-set $U(v)$ of v as follows:

$$D(v) = \{x \in V(G) : x \prec v\} \ \text{ and } \ U(v) = \{x \in V(G) : v \prec x\}.$$

For every vertex v, the collection of all the neighbours of v is denoted by $N(v)$.

Definition 3.3. *Let $A \subseteq V(G)$, and $\prec$ be a linear order of the vertex set of G. We define,*

$$\begin{aligned}\Gamma^*(G,\prec,A) &= \frac{1}{|V(G)|^3}\sum_{u\prec v}\big[|N(v)\cap A\cap D(u)| - |N(u)\cap A\cap D(u)|\big]_+\\ &+ \frac{1}{|V(G)|^3}\sum_{u\prec v}\big[|N(u)\cap A\cap U(v)| - |N(v)\cap A\cap U(v)|\big]_+,\end{aligned}$$

where

$$[x]_+ = \begin{cases} x & \text{if } x>0\\ 0 & \text{otherwise}\end{cases}.$$

We also define

$$\Gamma^*(G,\prec) = \max_{A\subseteq V(G)} \Gamma^*(G,\prec,A),$$

and

$$\Gamma^*(G) = \min_{\prec} \Gamma^*(G,\prec),$$

where the minimum is taken over all the linear orderings of $V(G)$.

Proposition 3.4. *A graph G is one-dimensional geometric if and only if $\Gamma^*(G) = 0$.*

Proof. Let G be a one-dimensional geometric graph, and A be an arbitrary subset of $V(G)$. Let $\prec$ be a linear ordering that satisfies Condition (5). Fix an arbitrary pair of vertices $u \prec v$ of G. By Condition (5), if z belongs to $N(v)\cap A\cap D(u)$ then z is adjacent to u as well. Thus $|N(v)\cap A\cap D(u)| \le |N(u)\cap A\cap D(u)|$. Similarly, one can show that $|N(u)\cap A\cap U(v)| \le |N(v)\cap A\cap U(v)|$, which implies that $\Gamma^*(G,\prec) = 0$. Thus $\Gamma^*(G) = 0$.

Conversely, let G be a graph such that $\Gamma^*(G) = 0$. Let $\prec$ be the linear order of $V(G)$ such that $\Gamma^*(G,\prec) = 0$. Let $u \prec v$ be an arbitrary pair of adjacent vertices of G, and take z so that $u \prec z \prec v$. Since $\Gamma^*(G,\prec,A) = 0$ for all $A \subseteq V(G)$, choosing $A = \{v\}$ gives that $1 = |N(u)\cap\{v\}\cap U(z)| \le |N(z)\cap\{v\}\cap U(z)|$. This implies that z is adjacent to v. Similarly, one can show that z is adjacent to u. Thus Condition (5) is satisfied for G, and G is a geometric graph. □

Next, we extend Condition (5) to functions in $\mathcal{W}_0$. The generalization is obtained by considering functions representing graphs, as introduced in the previous section. Let G be a one-dimensional geometric graph with a linear ordering $\prec$ of vertices that satisfies Condition (5). Let w_G be the function in $\mathcal{W}_0$ that represents G with respect to the labelling of $V(G)$ obtained from the linear ordering $\prec$. It follows that $w_G(x,z) = 1$ and $x \le y \le z$ imply that $w_G(x,y) = 1$ and $w_G(y,z) = 1$. We generalize this property as follows:

Definition 3.5. *A function $w \in \mathcal{W}$ is* diagonally increasing *if for every $x,y,z \in [0,1]$, we have*

1. $x \le y \le z \Rightarrow w(x,z) \le w(x,y)$,
2. $y \le z \le x \Rightarrow w(x,y) \le w(x,z)$.

A function w in $\mathcal{W}$ is diagonally increasing almost everywhere if there exists a diagonally increasing function w' which is equal to w almost everywhere.

Clearly, a graph is a one-dimensional geometric graph if and only if it has a function representative in $\mathcal{W}_0$ which is diagonally increasing. (Remember that we assume the function representative to have all blocks on the diagonal equal to 1.) Indeed, the function representative will be the function w_G where the vertices are ordered according to a linear ordering that satisfies Condition (5). More important is the connection between diagonally increasing functions and linear embeddings, as given below.

Proposition 3.6. *If $w \in \mathcal{W}_0$ is diagonally increasing, then the w-random graph $G(n, w)$ has a linear embedding.*

Proof. We may assume *w.l.o.g* that $w(x, x) = 1$ for all $x \in [0, 1]$. (Namely, if w and w' are equal almost everywhere, then $G(n, w) = G(n, w')$.) Consider $G(n, w)$. Its vertex set V is chosen *u.a.r* from $[0, 1]$. This constitutes a linear embedding. Namely, fix $\alpha \in [0, 1]$ and $v \in V$. Let $R = \{x \in \mathbb{R} : w(x, v) > \alpha\}$. Because w is diagonally increasing, R is an interval containing v, and hence convex. Since the probability that two vertices x and v are adjacent is equal to $w(x, v)$, the result follows. □

The converse is also true, in the sense that, if for all n, the w-random graph $G(n, w)$ has a linear embedding, then w^{ϕ} is diagonally increasing for some map $\phi \in \Phi$. (As explained in the previous section, graph sequences converge to equivalence classes of $\mathcal{W}_0$ given by transformations by such maps ϕ.) The proof follows as an easy corollary from the results presented later in this section.

Next, we introduce a parameter Γ which generalizes the graph parameter Γ^* to functions in $\mathcal{W}$. We will see that Γ identifies the diagonally increasing functions.

Definition 3.7. *Let $\mathcal{A}$ denote the collection of all measurable subsets of $[0, 1]$. Let w be a function in $\mathcal{W}$, and $A \in \mathcal{A}$. We define*

$$\begin{aligned}\Gamma(w, A) &= \int\int_{y<z} \Big[\int_{x \in A\cap[0,y]} (w(x, z) - w(x, y))\, dx\Big]_+ dydz \\ &+ \int\int_{y<z} \Big[\int_{x\in A\cap[z,1]} (w(x, y) - w(x, z))\, dx\Big]_+ dydz.\end{aligned}$$

Moreover, $\Gamma(w)$ is defined as

$$\Gamma(w) = \sup_{A\in\mathcal{A}} \Gamma(w, A),$$

where the supremum is taken over all the measurable subsets of $[0, 1]$.

It follows directly from the definitions that any function $w \in \mathcal{W}$ which is almost everywhere diagonally increasing has $\Gamma(w) = 0$. The converse also holds, as is stated in the following Proposition. The proof is technical, and is given in Section 5.

Proposition 3.8. *Let w be a function in $\mathcal{W}_0$. The function w is diagonally increasing almost everywhere if and only if $\Gamma(w) = 0$.*

A graph G can be represented as a function $w_G \in \mathcal{W}_0$, but it is not necessarily true that $\Gamma^*(G) = \Gamma(w_G)$, even when the representation w_G is obtained by using the ordering of the vertices that achieves $\Gamma^*(G)$. This is due to the fact that a set A which determines the value of $\Gamma(w)$ does not have to be consistent with the partition of $[0, 1]$ into n equal-sized parts on which w_G is defined. However, we can show that $\Gamma^*(G)$ and $\Gamma(w_G)$, computed using the same ordering of the vertices, are asymptotically equal. This follows directly from Theorem 5.1, which is stated and proved in Section 5.

Proposition 3.9. *Let G be a graph of size n, and let $\prec$ be a linear order of the vertices of G which achieves $\Gamma^*(G)$, i.e. $\Gamma^*(G) = \Gamma^*(G, \prec)$. Let w_G be the function in $\mathcal{W}_0$ that represents G with respect to the linear order $\prec$ of the vertices of G. Then*

$$\Gamma^*(G) = \Gamma(w_G) + \mathcal{O}(\frac{1}{n}).$$

Our final result is that the parameter Γ is continuous under the cut-norm. In particular, let $\{G_n\}$ be a graph sequence converging to a limit $w \in \mathcal{W}_0$, and let $\prec_n$ be appropriate labellings of vertices of G_n such that the functions in $\mathcal{W}_0$ representing G_n with respect to $\prec_n$ converge to w in cut-norm. Then $\Gamma^*(G_n, \prec_n)$ converges to $\Gamma(w)$. However, in order to study the limit of the sequence $\{\Gamma^*(G_n)\}$, we need to define the following parameter.

Definition 3.10. *Let w be a bounded function in $\mathcal{W}$. We define the new parameter $\tilde{\Gamma}$ to be*

$$\tilde{\Gamma}(w) = \inf_{\phi \in \Phi} \Gamma(w^{\phi}).$$

Note that $\tilde{(\Gamma)}$ extends the notion of Γ to the quotient space $\mathcal{W}/\Phi$.

Proposition 3.11. *Let $w \in \mathcal{W}_0$ be the limit of a convergent sequence $\{G_n\}_{n\in\mathbb{N}}$ of graphs with $|V(G_n)| \to \infty$. Then $\{\Gamma^*(G_n)\}_{n\in\mathbb{N}}$ converges to $\tilde{\Gamma}(w)$ as $n \to \infty$.*

This proposition follows as a corollary from Theorem 5.5, which is stated and proved in Section 5.

Specifically, this means that a graph sequence $\{G_n\}$ converges to a diagonally increasing function precisely when $\{\Gamma^*(G_n)\}$ goes to zero. If a graph sequence converges to a graph limit w, then the graphs in the sequence have approximately the same homomorphism densities as the w-random graph. Moreover, the w-random graph has a linear embedding. Therefore, the graphs in the sequence are structurally similar to those produced by a random graph process with a linear embedding, and thus their vertices have a natural interpretation as points in $\mathbb{R}$. Moreover, the optimal order for Γ^* gives the order in which points should be placed in $\mathbb{R}$. This justifies our assertion that Γ^* is a parameter that measures "embeddability" in $\mathbb{R}$.

4. Uniform linear embeddings and integer-valued graph limits

In a random graph with a spatial embedding, the probability of linking to a particular vertex v decreases with distance, but the rate of decrease may depend on the particular vertex v, and on the direction in which the distance increases. In this section, we consider the question when the embedding can be chosen to be uniform, so the probability of a link between two vertices depends only on the distance between them.

Definition 4.1. *A random graph on a vertex set V has a uniform spatial embedding into a metric space (S, d) if and only if there exists an injection $\pi : V \to S$ and a decreasing function $\delta_{pr} : [0, 1] \to \mathbb{R}^+ \cup \{0\}$ such that for every $a \in [0, 1]$ and for every $v \in V$, for all vertices $u \in V \setminus \{v\}$, $d(\pi(u), \pi(v)) < \delta_{pr}(a)$ if and only if $Prob(u \sim v) > a$.*

A function $w \in \mathcal{W}_0$ *has a uniform spatial embedding into a metric space* (S, d) *if there exists an injection* $\pi : [0,1] \to S$ *and a decreasing function* $f_{pr} : \mathbb{R}^+ \cup \{0\} \to [0,1]$ *such that for almost every* $x, y \in [0,1]$*,* $w(x,y) = f_{pr}(d(\pi(x), \pi(y)))$*.*

A uniform linear embedding is a uniform spatial embedding into $(\mathbb{R}, |\cdot|)$*.*

Clearly, for a given function $w \in \mathcal{W}_0$, the w-random graph $G(n, w)$ almost surely has a uniform spatial embedding precisely when w has a uniform spatial embedding. Moreover, the functions f_{pr} and δ_{pr} are each other's inverse, in the sense that, for all $a \in [0,1]$ and $x \in \mathbb{R}^+ \cup \{0\}$, $f_{pr}(x) < a$ if and only if $\delta_{pr}(a) < x$.

The function f_{pr} can be interpreted as the maximum distance where the link probability is above a given value. The function δ_{pr} gives the link probability as a function of distance. Clearly, a uniform spatial embedding is also a spatial embedding in the sense of Definition 3.1. In a spatial embedding, the embeddings of all vertices that have link probability greater than a to vertex v must be contained in a convex region containing v. In a uniform spatial embedding, this convex region must be a ball of radius $\delta_{pr}(a)$ around v.

In this section, we consider integer-valued graph limits, and answer the question when such graph limits have a uniform linear embedding. In particular, we take a closer look at diagonally increasing functions in $\mathcal{W}_0$ which only assume values zero or one. Consider such a function $w : [0,1] \to \{0,1\}$. From the properties given in Definition (3.5), we conclude that each point $x \in [0,1]$ can be associated with an interval $I_x = [\ell(x), r(x)]$, so that $w(x,y) = 1$ if and only if $y \in I_x$. (Note that the intervals could also be open or half open; we will assume that all intervals are closed, since this affects at most a set of measure zero.)

The functions ℓ and r are (not necessarily strictly) increasing. Namely, assume the contrary, so there exist x, y so that $x < y$ and $\ell(x) > \ell(y)$. Then there exists z so that $\ell(y) < z < \ell(x) \leq x < y$. So $z \in I_y$ and $z \notin I_x$, and thus $0 = w(x,z) < w(y,z) = 1$. This contradicts the fact that w is diagonally increasing. See Figure 1 for an example. Note that the discontinuity in ℓ corresponds to an interval where r is constant. Thus if r and ℓ are both strictly increasing, they are both continuous.

Since w is symmetric, for all $y < x$ we have that $y \geq \ell(x)$ if and only if $x \leq r(y)$. Thus, r completely determines w almost everywhere. Let $c = \inf\{x : r(x) = 1\}$, so $r(x) = 1$ on $(c, 1]$, and $r(x) < 1$ on $[0, c)$. See also Figure 1. If r is continuous and strictly increasing on $[0, c]$, then r can be realized as ℓ^{-1} on $[0, c]$, and 1 everywhere else.

If we shrink the intervals to half the size, or precisely, we define the associated interval $I^*_x = [x - \frac{x-\ell(x)}{2}, x + \frac{r(x)-x}{2}]$, then we have that $w(x,y) = 1$ if and only if $I^*_x \cap I^*_y \neq \emptyset$. Thus, w can be considered as an uncountably infinite interval graph. Moreover, since ℓ and r are increasing, we have that it is a proper interval graph. It is a folklore result that finite proper interval graphs can be represented as unit interval graphs (note that this also follows from Proposition 3.2.) In other words, there exists an embedding of the vertices in $\mathbb{R}$ so that vertices u and v are adjacent if and only if the distance between the embeddings of u and v is smaller than one.

We ask whether w satisfies a similar property. Precisely, does there exist an injective map $\pi : [0,1] \to \mathbb{R}$ so that for almost every x and y,

$$|\pi(x) - \pi(y)| < 1 \Rightarrow w(x,y) = 1, \quad \text{and } |\pi(x) - \pi(y)| > 1 \Rightarrow w(x,y) = 0? \tag{6}$$



FIGURE 1. An example of an integer-valued, diagonally increasing graph limit w. The grey area is where w equals 1, elsewhere w equals zero. The functions ℓ and r form the boundaries of the grey area.

Clearly, π is a uniform linear embedding in the sense of Definition 4.1. Moreover, any uniform linear embedding of w can be transformed in a map satisfying the property above by multiplication by a constant.

To see the relevance of this question, consider a sequence of one-dimensional random geometric graphs formed as follows. To generate G_n, n points are chosen from $\mathbb{R}$ according to a given distribution with a strictly increasing cumulative distribution function F (that is, the probability of choosing a member from the interval $[a, b)$ equals $F(b) - F(a)$). Points are made adjacent if and only if their distance is at most 1. This sequence converges to an integer-valued graph limit w_F given by

$$w_F(x, y) = 1 \text{ if and only if } |F^{-1}(x) - F^{-1}(y)| \leq 1.$$

Namely, the random graph G_n is exactly the w_F-random graph: choosing an element x uniformly at random from $[0, 1]$, and then placing it at $F^{-1}(x)$ is exactly the same process as choosing a point from $\mathbb{R}$ according to a distribution with *cdf* F.

The function w_F obtained from a random geometric graph process therefore has a uniform linear embedding given by F^{-1}. Conversely, if a uniform linear embedding π exists for a given integer-valued function w, and if π is strictly increasing, then the w-random graph is exactly the one dimensional random geometric graph where points are chosen from $\mathbb{R}$ according to a distribution with *cdf* $F = \pi^{-1}$.

Let us first introduce the following definitions.

Definition 4.2. *Let $w \in \mathcal{W}_0$ be diagonally increasing. Let $\ell(x)$ and $r(x)$ be so that $w(x, y) = 1$ if and only if $\ell(x) \leq y \leq r(x)$. We call w* consistent *if the functions ℓ and r are strictly increasing whenever they are* not *equal to zero or one. For any consistent function w, define*

$$\begin{aligned} L(w) &= \{z \in (0, 1] : \ell(z) = z\}, \text{ and} \\ R(w) &= \{z \in [0, 1) : r(z) = z\}. \end{aligned}$$

Any element z of L or R indicates a point where the functions ℓ and r meet the diagonal $x = y$. Thus, the intervals associated with points near z become infinitesimally small. Since, in the embedding, all intervals must have the same length, this can only be accommodated if the embedded values near z are allowed to go to infinity.

Theorem 4.3 below establishes when a uniform linear embedding exists, in the case that w is consistent. If w is consistent, then any element $z \in (0,1)$ for which $\ell(z) = z$ must also have $r(z) = z$. Therefore, z will be in $L(w) \cap R(w)$. The condition for existence of a uniform embedding, as given by the following theorem, is that $|L(w)|+|R(w)| \leq 2$. Thus, this means that either $L(w) \cup R(w) = \{z\}$, where $0 < z < 1$, or $L(w) \cup R(w) \subseteq \{0,1\}$.

We believe the results for the existence of uniform embeddings of $\{0,1\}$-valued functions can be extended to more general functions, and will explore this in future work. The proof of the theorem can be found in Section 5.

Theorem 4.3. *Let $w \in \mathcal{W}_0$ be diagonally increasing, integer-valued and consistent. Then w has a uniform linear embedding $\pi : (0,1) \to \mathbb{R}$ if and only if $|L(w)| + |R(w)| \leq 2$. A strictly increasing uniform linear embedding exists precisely when $L(w) \cup R(w) \subseteq \{0,1\}$. Moreover, $|\pi|$ is bounded if and only if $L(w) \cup R(w) = \emptyset$. In this case, π extends to $[0,1]$.*

We conclude with an example of a graph sequence which converges to an integer limit.

Example 1. Define a graph sequence $\{G_n\}$ as follows. Let $V(G_n) = \{z\} \cup X \cup Y$, where $X = \{x_1, \ldots, x_n\}$ and $Y = \{y_1, \ldots, y_n\}$. Vertex z is adjacent to x_1 and y_1, and no other vertices. No edges exist between X and Y. For all $i < j$, x_i is adjacent to x_j precisely when $j \leq 2i$, and y_i is adjacent to y_j precisely when $j \leq 2i$.

Clearly, each G_n is a one-dimensional geometric graph. In particular, if the vertices are ordered so that $y_i \prec y_j$ if and only if $i > j$, $x_i \prec x_j$ if and only if $i < j$, and $y_i \prec z$ and $z \prec x_i$ for all i, then G_n with this ordering satisfies Condition (5). The sequence $\{G_n\}$ with this ordering converges to an integer-valued, diagonally increasing graph limit w so that $w(x,y) = 1$ if and only if $y \in [\ell(x), r(x)]$, where

$$r(x) = \begin{cases} 1/4 + x/2 & \text{if } x \in [0,1/2] \\ 1/2 + 2(x - 1/2) & \text{if } x \in (1/2,3/4] \\ 1 & \text{if } x \in (3/4,1], \end{cases}$$

and $\ell(x)$ is defined accordingly.

Using the definitions from Theorem 4.3, we see that $L(w) = R(w) = \{1/2\}$. We can define a uniform linear embedding as follows:

$$\pi(x) = \begin{cases} \log_2(1/4 - x/2) & \text{if } x \in [0,1/2) \\ 0 & \text{if } x = 1/2 \\ -\log_2(x/2 - 1/4) & \text{if } x \in (1/2,1]. \end{cases}$$

It is straightforward to verify that π is a uniform linear embedding. For example, if $\{x,y\} \subseteq [0,1/2)$ and $x < y$, then $w(x,y) = 1$ precisely when $y \leq r(x) = 1/4 + x/2$. This occurs precisely when $1/2 - y \geq 1/4 - x/2$. By taking logarithms, this means that $\log_2(1/4 - y/2) + 1 \geq \log_2(1/4 - x/2)$ and thus $|\pi(x) - \pi(y)| \leq 1$.

Note that each graph G_n is connected, while the embedding has an isolated point corresponding to z, and two disjoint pieces corresponding to X and Y. Moreover, the embedding π does not respect the ordering of the vertices in $[0,1)$.

5. Proofs of the main results.

We start this section by introducing some notations which will be used later. Let $w \in \mathcal{W}$, and A and B be measurable subsets of $[0,1]$. We define $\tilde{w}(A,B)$ to be the average of w on $A \times B$, i.e.

$$\tilde{w}(A,B) = \frac{1}{\mu(A)\mu(B)} \int_{A\times B} w(x,y)dxdy,$$

where μ is the Lebesgue measure on $[0,1]$. Let n be a positive integer. For each $0 \le i \le n-1$, let $I_i = [\frac{i}{n}, \frac{i+1}{n}]$. We define the symmetric functions w_n, w_n^+, and w_n^- on $[0,1]^2$ as follows.

$$\begin{aligned}
w^n_{i,j} &= \tilde{w}(I_i, I_j) \text{ for } 0 \le i,j \le n-1, \\
w_n(x,y) &= w^n_{i,j} \text{ if } (x,y) \in I_i \times I_j, \\
w_n^-(x,y) &= \begin{cases} w^n_{i-1,j+1} & \text{if } (x,y) \in I_i \times I_j \ \&\ 1 \le i \le j \le n-2 \\ 0 & \text{if } (x,y) \in I_0 \times I_j \\ 0 & \text{if } (x,y) \in I_i \times I_{n-1}, \end{cases} \\
w_n^+(x,y) &= \begin{cases} w^n_{i+1,j-1} & \text{if } (x,y) \in I_i \times I_j \ \&\ i \le j-2 \\ 1 & \text{if } (x,y) \in I_i \times I_i \\ 1 & \text{if } (x,y) \in I_i \times I_{i+1}. \end{cases}
\end{aligned}$$

Let A and B be subsets of $[0,1]$. We say $A \le B$ if every a in A is smaller than or equal to every b in B.

We now give the proof of Proposition 3.8, which states that a function $w \in \mathcal{W}_0$ is diagonally increasing almost everywhere if and only if $\Gamma(w) = 0$. This proof is inspired by the proof of Lemma 4.6 of [3]. However, we include the proof to make the paper self-contained.

Proof of Proposition 3.8. Clearly, if w is diagonally increasing almost everywhere then $\Gamma(w) = 0$. To prove the other direction, let w be a function in $\mathcal{W}_0$ with $\Gamma(w) = 0$. Let A, B, and C be subsets of $[0,1]$ such that $C \le A \le B$. Since $\Gamma(w) = 0$, for almost every $y \in A$ and almost every $z \in B$,

$$\int_{x\in C} w(x,z)dx \le \int_{x \in C} w(x,y)dx. \tag{7}$$

Taking repeated integrals of both sides of Equation (7) over A and then B and then dividing by $\mu(A)$, we conclude that

$$\int_{C\times B} w(x,z)dxdz \le \frac{\mu(B)}{\mu(A)} \int_{C\times A} w(x,y)dxdy. \tag{8}$$

Similarly, one can show that for subsets A, B, and C of $[0,1]$ with $A \le B \le C$, we have

$$\int_{A\times C} w(x,y)dydx \le \frac{\mu(A)}{\mu(B)} \int_{B\times C} w(x,z)dzdx. \tag{9}$$

Applying the above inequalities to the sets I_i, we have that for every $(x,y) \in [0,1]^2$, $w_n^-(x,y) \le w_n(x,y) \le w_n^+(x,y)$. Now let A and B be subsets of $[0,1]$. From

Equations (8) and (9) it follows that, if $0 \le i \le j-2 \le n-3$, then

$$\begin{aligned}\int_{(A\cap I_i)\times(B\cap I_j)} w(x,y)dxdy &\le \frac{\mu(B\cap I_j)}{\mu(I_{j-1})}\int_{(A\cap I_i)\times I_{j-1}} w(x,y)dxdy\\ &\le \frac{\mu(A\cap I_i)\mu(B\cap I_j)}{\mu(I_{i+1})\mu(I_{j-1})}\int_{I_{i+1}\times I_{j-1}} w(x,y)dxdy\\ &= \mu(A\cap I_i)\mu(B\cap I_j)w^n_{i+1,j-1}.\end{aligned}$$

Using the fact that w is symmetric, we obtain

$$\begin{aligned}&\int_{A\times B} w(x,y)dxdy\\
=\ &\sum_{i,j=0}^{n-1}\int_{(A\cap I_i)\times(B\cap I_j)} w(x,y)dxdy\\
=\ &\sum_{0\le i\le j-2\le n-3}\int_{(A\cap I_i)\times(B\cap I_j)} w(x,y)dxdy+\int_{(B\cap I_i)\times(A\cap I_j)} w(x,y)dxdy\\
&\quad+\sum_{i=0}^{n-2}\int_{(A\cap I_i)\times(B\cap I_{i+1})} w(x,y)dxdy+\int_{(B\cap I_i)\times(A\cap I_{i+1})} w(x,y)dxdy\\
&\quad+\sum_{i=0}^{n-1}\int_{(A\cap I_i)\times(B\cap I_i)} w(x,y)dxdy\\
\le\ &\sum_{0\le i\le j-2\le n-3}(\mu(A\cap I_i)\mu(B\cap I_j)+\mu(B\cap I_i)\mu(A\cap I_j))w^n_{i+1,j-1}\\
&\quad+\sum_{i=0}^{n-2}\mu(A\cap I_i)\mu(B\cap I_{i+1})+\mu(B\cap I_i)\mu(A\cap I_{i+1})\\
&\quad+\sum_{i=0}^{n-1}\mu(A\cap I_i)\mu(B\cap I_i)\\
=\ &\sum_{0\le i\le j-2\le n-3}\int_{(A\cap I_i)\times(B\cap I_j)} w_n^+(x,y)dxdy+\int_{(B\cap I_i)\times(A\cap I_j)} w_n^+(x,y)dxdy\\
&\quad+\sum_{i=0}^{n-2}\int_{(A\cap I_i)\times(B\cap I_{i+1})} w_n^+(x,y)dxdy+\int_{(B\cap I_i)\times(A\cap I_{i+1})} w_n^+(x,y)dxdy\\
&\quad+\sum_{i=0}^{n-1}\int_{(A\cap I_i)\times(B\cap I_i)} w_n^+(x,y)dxdy\\
=\ &\int_{A\times B} w_n^+(x,y)dxdy.\end{aligned}$$

Moreover, since measurable subsets of $[0,1]^2$ can be approximated in measure by finite union of disjoint rectangles, we get

$$\int_E w(x,y)dxdy \le \int_E w_n^+(x,y)dxdy,$$

for every measurable subset E of $[0,1]^2$. Thus, $w \le w_n^+$ (and similarly $w_n^- \le w$) almost everywhere in $[0,1]^2$. This, together with the definitions of w_n^+ and w_n^-, implies that

$$\|w-w_n\|_1 \leq \|w_n^+ - w_n^-\|_1 = \int_{[0,1]^2}(w_n^+ - w_n^-)(x,y)dxdy \leq \frac{8}{n}.$$

Using the Borel-Cantelli lemma, we conclude that the sequence $\{w_{2^n}\}_{n\in\mathbb{N}}$ converges to w almost everywhere in $[0,1]^2$, i.e. $\psi := \limsup_{n\in\mathbb{N}} w_{2^n} = w$ almost everywhere. Finally, by Equations (8) and (9), each w_n is a diagonally increasing function. Therefore, ψ is diagonally increasing as well. □

Next is the proof of the fact that for any graph G, the difference between the graph parameter $\Gamma^*(G)$ and the associated function parameter $\Gamma(w_G)$ is bounded by a function of n, the size of G, which goes to zero as n grows large. Since w_G is a function constant on the sets $I_i \times I_j$, this result follows directly from the following theorem. Recall from Definition 3.7 that $\mathcal{A}$ denotes the collection of all measurable subsets of $[0,1]$.

Theorem 5.1. *Let $n \in \mathbb{N}$. Let $w \in \mathcal{W}$ be a function bounded by c, which is measurable with respect to the product algebra $\mathcal{A}_n^* \times \mathcal{A}_n^*$, where the algebra $\mathcal{A}_n^*$ is generated by the intervals $\{I_i : 0 \leq i \leq n-1\}$. Then $\Gamma(w) = \sup_{A\in\mathcal{A}}\Gamma(w,A) = \max_{A\in\mathcal{A}_n^*}\Gamma(w,A) + \mathcal{O}(\frac{c}{n})$.*

Proof. Let $n \in \mathbb{N}$ and $w \in \mathcal{W}$ be as above. Note that w is constant on the rectangles $I_i \times I_j$, since it is measurable with respect to the product algebra $\mathcal{A}_n^* \times \mathcal{A}_n^*$. For each $i,j \in \{0,\ldots,n-1\}$, let $w(x,y) = a_{ij}$ whenever $(x,y) \in I_i \times I_j$. Fix $A \in \mathcal{A}$, and let $\beta_k = \mu(A \cap I_k)$ for every $0 \leq k \leq n-1$. The expression for $\Gamma(w,A)$ as given in Definition 3.7 can now be simplified.

Consider $y < z$ so that $y \in I_i$ and $z \in I_j$. If $i = j$, then for all x, $w(x,z) = w(x,y)$, so $\int_{x\in A\cap[0,y]}(w(x,z)-w(x,y))\,dx\big]_+ = 0$. If $0 \leq i < j \leq n-1$, then

$$\begin{aligned}
&\Big[\int_{x\in A\cap[0,y]}(w(x,z)-w(x,y))\,dx\Big]_+ \\
= \;& \Big[\sum_{k=0}^{i-1}\int_{A\cap I_k}(a_{kj}-a_{ki})dx + \int_{A\cap I_i\cap[0,y]}(a_{ij}-a_{ii})dx\Big]_+ \\
= \;& \Big[\sum_{k=0}^{i-1}\mu(A\cap I_k)(a_{kj}-a_{ki}) + \mu(A\cap I_i\cap[0,y])(a_{ij}-a_{ii})\Big]_+ \\
\leq \;& \left(\Big[\sum_{k=0}^{i-1}\beta_k(a_{kj}-a_{ki})\Big]_+ + \frac{2c}{n}\right).
\end{aligned}$$

In the last step, we use the inequality $[x+y]_+ \leq [x]_+ + [y]_+$, and the fact that w is bounded by c, so $|\mu(A\cap I_i\cap[0,y])(a_{ij}-a_{ii})|$ is at most $\frac{2c}{n}$.

Similarly, we have that

$$\begin{aligned}
& \Big[\int_{x\in A\cap[z,1]} (w(x,y)-w(x,z))\,dx\Big]_+ \\
= & \Big[\sum_{k=j+1}^{n-1} \mu(A\cap I_k)(a_{ki}-a_{kj}) + \mu(A\cap I_j\cap[z,1])(a_{ji}-a_{jj})\Big]_+ \\
\leq & \left(\Big[\sum_{k=j+1}^{n-1} \beta_k(a_{ki}-a_{kj})\Big]_+ + \frac{2c}{n}\right).
\end{aligned}$$

Using this, we can bound $\Gamma(w,A)$:

$$\begin{aligned}
\Gamma(w,A) \leq & \sum_{0\leq i<j\leq n-1} \int_{y\in I_i}\int_{z\in I_j} \left(\Big[\sum_{k=0}^{i-1}\beta_k(a_{kj}-a_{ki})\Big]_+ + \frac{2c}{n}\right) dydz \\
+ & \sum_{0\leq i<j\leq n-1} \int_{y\in I_i}\int_{z\in I_j} \left(\Big[\sum_{k=j+1}^{n-1}\beta_k(a_{ki}-a_{kj})\Big]_+ + \frac{2c}{n}\right) dydz \\
= & \sum_{0\leq i<j\leq n-1} \frac{1}{n^2}\Big[\sum_{k=0}^{i-1}\beta_k(a_{kj}-a_{ki})\Big]_+ + \frac{c(n-1)}{n^2} \\
+ & \sum_{0\leq i<j\leq n-1} \frac{1}{n^2}\Big[\sum_{k=j+1}^{n-1}\beta_k(a_{ki}-a_{kj})\Big]_+ + \frac{c(n-1)}{n^2}.
\end{aligned}$$

Now define,

$$\begin{aligned}
g_w(A) &= g_w(\beta_0,\ldots,\beta_{n-1}) \\
&= \sum_{0\leq i<j\leq n-1} \frac{1}{n^2}\left(\Big[\sum_{k=0}^{i-1}\beta_k(a_{kj}-a_{ki})\Big]_+ + \Big[\sum_{k=j+1}^{n-1}\beta_k(a_{ki}-a_{kj})\Big]_+\right).
\end{aligned}$$

Thus,

$$\Gamma(w,A) \leq g_w(A) + \frac{2c(n-1)}{n^2} \leq g_w(A) + \frac{2c}{n}. \tag{10}$$

Similarly, one can use the inequality $[x+y]_+ \geq [x]_+ - |y|$ to show that

$$\Gamma(w,A) \geq g_w(A) - \frac{2c}{n}. \tag{11}$$

Since $x\mapsto [x]_+$ is a convex function, g_w is the sum of convex functions, and therefore is itself also convex. Moreover, since $\beta_k\in[0,\frac{1}{n}]$, the function g_w achieves its maximum when each of the coefficients β_k is either 0 or $\frac{1}{n}$. Since $\beta_k=\mu(A\cap I_k)$, this implies that the maximum is achieved when, for each k, either A contains I_k, or is disjoint from I_k. Hence, $\sup_{A\in\mathcal{A}} g_w(A) = \max_{A\in\mathcal{A}_n^*} g_w(A)$.

Let $A' \in \mathcal{A}_n^*$ be such that $\max_{A\in\mathcal{A}_n^*} g_w(A) = g_w(A')$. Then, by Equation (10) and (11) we have,

$$\begin{aligned} \sup_{A\in\mathcal{A}} \Gamma(w,A) &\leq \sup_{A\in\mathcal{A}} g_w(A) + \frac{2c}{n} = \max_{A\in\mathcal{A}_n^*} g_w(A) + \frac{2c}{n} \\ &= g_w(A') + \frac{2c}{n} \leq \Gamma(w,A') + \frac{4c}{n} \\ &\leq \max_{A\in\mathcal{A}^*} \Gamma(w,A) + \frac{4c}{n}. \end{aligned} \tag{12}$$

On the other hand, we clearly have

$$\max_{A\in\mathcal{A}_n^*} \Gamma(w,A) \leq \sup_{A\in\mathcal{A}} \Gamma(w,A). \tag{13}$$

Finally Equations (12) and (13) imply that

$$|\sup_{A\in\mathcal{A}} \Gamma(w,A) - \max_{A\in\mathcal{A}_n^*} \Gamma(w,A)| \leq \frac{4c}{n}.$$

□

Corollary 5.2. *Let G be a graph with n vertices, and w_G be the function in $\mathcal{W}_0$ that represents G with respect to a linear ordering $\prec$ of the vertices of G. Then*

$$\Gamma^*(G,\prec) = \Gamma(w_G) + \mathcal{O}(\frac{1}{n}).$$

Proof. Let $A \in \mathcal{A}_n^*$, and define $\tilde{A} = \{0 \leq i \leq n-1 : I_i \subseteq A\}$. From the proof of Theorem 5.1, it is easy to observe that $\Gamma^*(G,\prec,\tilde{A}) = g_{w_G}(A)$, and $g_{w_G}(A) - \frac{2}{n} \leq \Gamma(w_G,A) \leq g_{w_G}(A) + \frac{2}{n}$. Thus,

$$\max_{A\in\mathcal{A}_n^*} \Gamma^*(G,\prec,\tilde{A}) - \frac{2}{n} \leq \max_{A\in\mathcal{A}_n^*} \Gamma(w_G,A) \leq \max_{A\in\mathcal{A}_n^*} \Gamma^*(G,\prec,\tilde{A}) + \frac{2}{n}.$$

Using Theorem 5.1, we conclude that $|\Gamma^*(G,\prec) - \Gamma(w_G)| \leq \frac{6}{n}$, and we are done. □

Finally, we prove that the function parameter Γ is continuous under the cut-norm. This is a direct consequence of Theorem 5.5. Let us first present the following lemma.

Lemma 5.3. *Let $w : [0,1]^2 \to [-2,2]$ be a measurable function with $\|w\|_\square \leq \frac{1}{2}$. Then $\|w\chi\|_\square \leq 2\sqrt{\|w\|_\square}$, where*

$$\chi(x,y) = \begin{cases} 1 & x \leq y \\ 0 & \textit{otherwise} \end{cases}.$$

Proof. Let $\Omega = \{(x,y) : 0 \leq x \leq y \leq 1\}$ denote the subset of points above the diagonal in $[0,1]^2$. Define $k = \lceil \frac{1}{\sqrt{\|w\|_\square}} \rceil$, which is at least two by our condition on w. Now, we can decompose Ω into $k-1$ rectangles and k triangles as shown in Figure 5. Precisely, the i-th rectangle has width $\frac{1}{k}$ and ranges from $y = \frac{i}{k}$ to $y = 1$, and each triangle has base and height equal to $\frac{1}{k}$. By the definition of the cut norm, the integral of w over each of the rectangles is at most $||w||_\square$, in absolute value. Also, each of the triangles has measure $\frac{1}{2k^2}$, and there are k triangles in total. Since $|w|$ is bounded by 2, the integral of w over the triangles is at most $\frac{1}{2k^2}(2)(k) = \frac{1}{k}$, in absolute value. Therefore, we have



FIGURE 2. The decomposition of the set of points above the diagonal as used in the proof of Lemma 5.3

$$\begin{aligned}\left|\int_0^1\int_0^1 w\chi(x,y)d_xd_y\right| &\leq \frac{1}{k}+(k-1)\|w\|_\square\\ &\leq \sqrt{\|w\|_\square}+(\frac{1}{\sqrt{\|w\|_\square}})\|w\|_\square = 2\sqrt{\|w\|_\square}.\end{aligned}$$

For arbitrary subsets A and B of $[0,1]$, let $\chi_{A\times B}$ denote the characteristic function of the subset $A\times B$ of $[0,1]^2$. Clearly $|\int_0^1\int_0^1 w\chi_{A\times B}\chi(x,y)d_xd_y| \leq 2\sqrt{\|w\chi_{A\times B}\|_\square} \leq 2\sqrt{\|w\|_\square}$, which proves that $\|w\chi\|_\square \leq 2\sqrt{\|w\|_\square}$. □

Lemma 5.4. *Let w_1 and w_2 be elements of $\mathcal{W}_0$ with cut-norms bounded by $\frac{1}{4}$. Then $|\Gamma(w_1)-\Gamma(w_2)| \leq 2\|w_1-w_2\|_\square + 2\sqrt{\|w_1-w_2\|_\square}$.*

Proof. From the definition of Γ, let

$$\begin{aligned}\Gamma_1(w,A) &= \int\int_{y<z}\left[\int_{x\in A\cap[0,y]}(w(x,z)-w(x,y))\,dx\right]_+ dydz,\\ \Gamma_2(w,A) &= \int\int_{y<z}\left[\int_{x\in A\cap[z,1]}(w(x,y)-w(x,z))\,dx\right]_+ dydz.\end{aligned}$$

Fix a measurable set $A \in \mathcal{A}$. Using again the inequality $[x+y]_+ \le [x]_+ + [y]_+$, we obtain that

$$\begin{aligned}
\Gamma_1(w_1, A) &= \int\int_{y<z} \left[\int_{x \in A \cap [0,y]} (w_1(x,z) - w_1(x,y))\, dx \right]_+ dydz \\
&\le \int\int_{y<z} \left[\int_{x \in A \cap [0,y]} (w_1(x,z) - w_2(x,z))\, dx \right]_+ dydz \\
&+ \int\int_{y<z} \left[\int_{x \in A \cap [0,y]} (w_2(x,z) - w_2(x,y))\, dx \right]_+ dydz \\
&+ \int\int_{y<z} \left[\int_{x \in A \cap [0,y]} (w_2(x,y) - w_1(x,y))\, dx \right]_+ dydz.
\end{aligned}$$

There exists $y_0, z_0 \in [0,1]$ such that

$$\begin{aligned}
\Gamma_1(w_1, A) &\le \int_{y_0<z} \left[\int_{x \in A \cap [0,y_0]} (w_1(x,z) - w_2(x,z))\, dx \right]_+ dz \\
&+ \Gamma_1(w_2, A) \\
&+ \int_{y<z_0} \left[\int_{x \in A \cap [0,y]} (w_2(x,y) - w_1(x,y))\, dx \right]_+ dy \\
&= \int_{z \in T_1} \int_{x \in A \cap [0,y_0]} (w_1(x,z) - w_2(x,z))\, dxdz \\
&+ \Gamma_1(w_2, A) \\
&+ \int_{y \in T_2} \int_{x \in A \cap [0,y]} (w_2(x,y) - w_1(x,y))\, dxdy,
\end{aligned}$$

where T_1 and T_2 are the appropriate sets of points which make the associated expressions positive. From the definition of the cut-norm, it then follows that

$$\Gamma_1(w_1, A) - \Gamma_1(w_2, A) \le \|w_1 - w_2\|_\square + \|(w_1 - w_2)\chi\|_\square.$$

Similarly, by switching w_1 and w_2, we get $\Gamma_1(w_2, A) - \Gamma_1(w_1, A) \le \|w_1 - w_2\|_\square + \|(w_1 - w_2)\chi\|_\square$, which implies that

$$|\Gamma_1(w_1, A) - \Gamma_1(w_2, A)| \le \|w_1 - w_2\|_\square + \|(w_1 - w_2)\chi\|_\square$$

holds for every subset A. Moreover, one can prove the analogue result for Γ_2. Thus,

$$\begin{aligned}
|\Gamma(w_1, A) - \Gamma(w_2, A)| &\le |\Gamma_1(w_1, A) - \Gamma_1(w_2, A)| + |\Gamma_2(w_1, A) - \Gamma_2(w_2, A)| \\
&\le 2\|w_1 - w_2\|_\square + 2\|(w_1 - w_2)\chi\|_\square.
\end{aligned}$$

Let $\epsilon > 0$ be given. There exists $A \subseteq [0,1]$ such that $\Gamma(w_1) \le \Gamma(w_1, A) + \epsilon$. By the definition of Γ, $\Gamma(w_2, A) \le \Gamma(w_2)$. Thus,

$$\Gamma(w_1) - \Gamma(w_2) \le \Gamma(w_1, A) + \epsilon - \Gamma(w_2, A) \le 2\|w_1 - w_2\|_\square + 2\|(w_1 - w_2)\chi\|_\square + \epsilon,$$

which, by switching w_1 and w_2, implies that

$$|\Gamma(w_1) - \Gamma(w_2)| \le 2\|w_1 - w_2\|_\square + 2\|(w_1 - w_2)\chi\|_\square.$$

This fact, together with Lemma 5.3, finishes the proof. □

Recall the definition of $\tilde{\Gamma}$ as given in Definition 3.10. Recall also that Φ is the set of all invertible maps $\phi : [0,1] \to [0,1]$ such that both ϕ and is inverse are measure preserving. The lemmas above lead to the following theorem, which establishes the continuity of the parameter $\tilde{\Gamma}$ on the space $\mathcal{W}_0$ with the cut distance $\delta_\square$.

Theorem 5.5. *Let $w \in \mathcal{W}_0$ be the limit of a $\delta_\square$-convergent sequence $\{w_n\}_{n\in\mathbb{N}}$ of functions in $\mathcal{W}_0$. Then $\{\tilde{\Gamma}(w_n)\}_{n\in\mathbb{N}}$ converges to $\tilde{\Gamma}(w)$ as $n \to \infty$.*

Proof. Since $\tilde{\Gamma}(w) = \inf_{\phi\in\Phi} \Gamma(w^\phi)$, for each positive integer m there must exist an invertible map $\phi_m \in \Phi$ such that $|\Gamma(w^{\phi_m}) - \tilde{\Gamma}(w)| \leq \frac{1}{m}$. Fix such a sequence of maps $\{\phi_m\}_{m=1}^\infty$.

Fix $m \in \mathbb{N}$. Then

$$\delta_\square(w_n, w^{\phi_m}) = \delta_\square(w_n, w) \to 0,$$

as n goes to infinity. By Theorem 2.3 of [9], this convergence implies that there exist maps $\psi_n \in \Phi$ such that

$$\|w_n^{\psi_n} - w^{\phi_m}\|_\square \to 0 \ \text{ as } n \to \infty.$$

By Lemma 5.4 we have,

$$\Gamma(w_n^{\psi_n}) \to \Gamma(w^{\phi_m})$$

Thus, for every $m \in \mathbb{N}$,

$$\limsup_{n\in\mathbb{N}} \tilde{\Gamma}(w_n) \leq \limsup_{n\in\mathbb{N}} \Gamma(w_n^{\psi_n}) = \Gamma(w^{\phi_m}) \leq \tilde{\Gamma}(w) + \frac{1}{m},$$

which implies that $\limsup_{n\in\mathbb{N}} \tilde{\Gamma}(w_n) \leq \tilde{\Gamma}(w)$.

To prove the other inequality, let $\gamma := \liminf_{n\in\mathbb{N}} \tilde{\Gamma}(w_n)$, and recall that, by assumption, $\delta_\square(w_n, w) \to 0$ as $n \to \infty$. Fix $0 < \epsilon < 1$, and let $n \in \mathbb{N}$ be chosen such that it satisfies

$$\delta_\square(w_n, w) < \frac{\epsilon^2}{12^2}, \text{ and } |\tilde{\Gamma}(w_n) - \gamma| < \frac{\epsilon}{3}.$$

In addition, let $\psi_n \in \Phi$ be such that $|\Gamma(w_n^{\psi_n}) - \tilde{\Gamma}(w_n)| < \epsilon/3$. By definition of the $\delta_\square$-distance, there exists $\phi \in \Phi$ such that $\|w_n^{\psi_n} - w^\phi\|_\square < \frac{\epsilon^2}{12^2}$ and thus, by Lemma 5.4, $|\Gamma(w_n^{\psi_n}) - \Gamma(w^\phi)| \leq 4\sqrt{||w_n^{\psi_n} - w^\phi||_\square} < \epsilon/3$. Thus,

$$\begin{aligned} |\Gamma(w^\phi) - \gamma| &\leq |\Gamma(w^\phi) - \Gamma(w_n^{\psi_n})| + |\Gamma(w_n^{\psi_n}) - \tilde{\Gamma}(w_n)| + |\tilde{\Gamma}(w_n) - \gamma| \\ &< \epsilon. \end{aligned}$$

Therefore, for every $0 < \epsilon < 1$, $\tilde{\Gamma}(w) \leq \Gamma(w^\phi) \leq \liminf_{n\in\mathbb{N}} \tilde{\Gamma}(w_n) + \epsilon$. Combining this with the lower bound, we get

$$\limsup_{n\in\mathbb{N}} \tilde{\Gamma}(w_n) \leq \tilde{\Gamma}(w) \leq \liminf_{n\in\mathbb{N}} \tilde{\Gamma}(w_n),$$

which implies that $\lim_{n\to\infty} \tilde{\Gamma}(w_n) = \tilde{\Gamma}(w)$.

□

Corollary 5.6. *Let $w \in \mathcal{W}_0$ be the limit of a convergent sequence $\{G_n\}_{n\in\mathbb{N}}$ of graphs with $|V(G_n)| \to \infty$. Then $\{\Gamma^*(G_n)\}_{n\in\mathbb{N}}$ converges to $\tilde{\Gamma}(w)$ as $n \to \infty$.*

Proof. By Corollary 5.2, for any increasing graph sequence $\{G_n\}$, the sequence $\Gamma^*(G_n)$ converges if and only if the sequence $\Gamma(w_{G_n})$ converges, where w_{G_n} is the step function representing G_n with respect to an ordering that is optimal for Γ^*. Moreover, if they exist, these limits are equal. Applying Theorem 5.5, the result follows. □

We conclude with the proof of Theorem 4.3 about the existence of a uniform embedding.

Proof of Theorem 4.3. Let r and ℓ be the functions defined on $[0,1]$ as in Definition 4.2. That is, for all pairs (x,y), $w(x,y)=1$ if $\ell(x)\leq y\leq r(x)$, and 0 otherwise. Let $c=\ell(1)$. The statement of the theorem assumes that w is consistent, so r and ℓ are both continuous. Also, r is strictly increasing on $[0,c]$, and ℓ restricted to $[r(0),1]$ equals the inverse of r restricted to $[0,c]$. It is straightforward to verify that for all $x<y$, $y\leq r(x)$ if and only if $x\geq \ell(y)$. Define a sequence $\{x_i\}_{i\geq 0}$ recursively by taking

$$x_0=0, \text{ and } x_{i+1}=r(x_i) \text{ for all } i\geq 0. \tag{14}$$

Since $\{x_i\}$ is increasing and bounded by 1, it converges to a limit $a\leq 1$. We will use the facts listed in the following claim.

Claim 5.7. *Let x_i and a be as above. Then,*

(i) *If $x_i<a$ for every $i\geq 0$, then $\ell(a)=a$, so $a\in L(w)$.*
(ii) *If there exists $k\geq 0$ such that $x_i=a$ for every $i\geq k$, then $a\in\{0,1\}$.*
(iii) $(0,a)\cap L(w)=\emptyset$.
(iv) $a\in L(w)\cup R(w)\cup\{0,1\}$.

Proof of claim: By definition, $\ell(a)\leq a$. On the other hand, for every $i\geq 0$, $\ell(a)>x_i$, since $x_{i+1}=r(x_i)<a$. Thus $\ell(a)\geq a$, which proves (i).

Suppose that there exists $k\geq 0$ such that $x_i=a$ for every $i\geq k$, and $x_i<a$ for $i<k$. Obviously, $r(a)=a$. If $k=0$, then $a=x_0=0$ and $a\in R(w)$. Assume $k>0$, then $r(x_{k-1})=x_k=a=r(a)$, while $x_{k-1}<a$. Since r is strictly increasing on $[0,c]$, where $c=\ell(1)$, this implies that $a>c$. So $r(a)=1$, and thus $a=1$.

To prove (iii), suppose there exists $z\in(0,a)\cap L(w)$. Since $(0,a)\subseteq\cup_{i\geq 0}(x_i,x_{i+1}]$, there exists an index i such that $x_i<z\leq x_{i+1}$. Now $\ell(z)=z$, so $\ell(z)>x_i$ and thus $z>r(x_i)=x_{i+1}$, which is a contradiction.

Part (iv) directly follows from (i) and (ii) and the definitions of $L(w)$ and $R(w)$.

Let us now define a uniform embedding π recursively on each of the intervals (x_i,x_{i+1}) as follows. Let $\pi(x_i)=i$ for all i, and

$$\pi(x)=\begin{cases}\frac{x}{x_1} & \text{if } x\in(x_0,x_1),\\ \pi(\ell(x))+1 & \text{if } x\in(x_i,x_{i+1}) \text{ for some } i\geq 1.\end{cases} \tag{15}$$

If $x\in(x_i,x_{i+1}]$ for some $i>0$, then $\ell(x)\in(x_{i-1},x_i]$. Namely, assume, by contradiction, that $\ell(x)>x_i$. Then $x>r(x_i)=x_{i+1}$, which contradicts the assumption. Moreover, since ℓ is strictly increasing on $[r(0),1]$ and $x_1=r(0)$, we have that $\ell(x)>\ell(x_i)=x_{i-1}$. Thus, π is well-defined on $[0,a)$. Moreover, since the function $\frac{x}{x_1}$ is strictly increasing, and ℓ is strictly increasing on $[x_1,1]$, π is strictly increasing on $[0,a)$. Clearly, π is an increasing uniform linear embedding of $w|_{[0,a)\times[0,a)}$ into $\mathbb{R}^{\geq 0}$, i.e. every pair $x,y\in[0,a)$ satisfies Equation (6). If $x_k=a$ for some index $k\in\mathbb{N}$, then the embedding extends to a, so π is an embedding of $w|_{[0,a]\times[0,a]}$. Moreover, in this case π is bounded above by k.

Assume that $|L(w)| + |R(w)| \leq 2$. Recall that this means that either $L(w) \cup R(w) = \{z\}$, where $0 < z < 1$, or $L(w) \cup R(w) \subseteq \{0, 1\}$. Consider the following cases:

Case 1: Suppose that $L(w) \cup R(w) \subseteq \{1\}$. By definition, this implies that $R(w) = \emptyset$, so $r(0) > 0$, and thus $a \geq x_1 > 0$. Therefore, $a = 1$, and π is a uniform linear embedding from $[0, 1)$ into $\mathbb{R}^{\geq 0}$. By (i) of the claim, if $x_i < 1$ for every $i \geq 0$, then $1 \in L(w)$, and π is unbounded. If $L(w) = \emptyset$, then there must exist an integer k with $x_k = 1$. In this case, π is a bounded uniform linear embedding from $[0, 1]$ into $\mathbb{R}^{\geq 0}$.

Case 2: Assume $L(w) \cup R(w) = \{0\}$, so $R(w) = \{0\}$ and $L(w) = \emptyset$. Let $\phi : [0, 1] \to [0, 1]$ be defined as $\phi(x) = 1 - x$. Clearly, ϕ is a one-to-one, onto, measure-preserving map which reverses the order on $[0, 1]$, and w^{ϕ} is a diagonally increasing integer-valued element of $\mathcal{W}_0$. It is easy to see that $z \in L(w)$ (respectively $R(w)$) if and only if $\phi^{-1}(z) \in R(w^{\phi})$ (respectively $L(w^{\phi})$). Thus, $L(w^{\phi}) = \{1\}$ and $R(w^{\phi}) = \emptyset$. Thus w^{ϕ} satisfies the conditions of Case 1, and there exists an (unbounded) uniform linear embedding π of w^{ϕ} into $\mathbb{R}^{\geq 0}$ that satisfies Equation (6). Clearly, $\pi \circ \phi^{-1} : (0, 1] \to \mathbb{R}^{\geq 0}$ is a uniform linear embedding of w.

Case 3: Assume $L(w) \cup R(w) \subseteq (0, 1)$. Let $\{x_i\}$ be the sequence defined in Equation (14) and let a be its limit. By (iv) of the claim and the assumption of this case, we must have that $L(w) = R(w) = \{a\}$ and $0 < a < 1$. In fact, since ℓ is strictly increasing on $[r(0), 1]$ and $a \geq x_1 = r(0)$, we have that $a = \ell(a) < \ell(1) = c$. Since r is strictly increasing on $[0, c]$, we can argue by induction that $x_i < a$ for all $i \geq 0$, and hence the sequence $\{x_i\}_{i \in \mathbb{N}}$ is infinite. Namely, assume $x_{i-1} < a$, then $x_i = r(x_{i-1}) < r(a) = a$.

Thus, the map $\pi : [0, a) \to \mathbb{R}^{\geq 0}$ defined as in Equation (15) forms an (unbounded) uniform linear embedding of $w|_{[0,a)\times[0,a)}$.

To complete the proof, note that the function $w|_{[a,1]\times[a,1]}$ behaves as described in Case 2, and one can apply similar methods to embed the interval $(a, 1]$ into $\mathbb{R}^{\leq -2}$. Namely, define a second sequence $\{y_i\}$ by $y_0 = 1$, $y_{i+1} = \ell(y_i)$ for all $i \geq 0$. This will lead to an infinite sequence with limit a. Now set $\pi(y_i) = -2 - i$ for all i, set $\pi(y) = -2 - \frac{y - y_0}{y_1 - y_0}$ for $y \in (y_0, y_1)$, and $\pi(r(y)) - 1$ for $y \in (y_i, y_{i+1})$, $i \geq 1$. Finally, let $\pi(a) = -1$. Since a has no neighbours in $[0, a)$, this is a uniform linear embedding. Note that, in this case π is not an increasing map, so the ordering of the embedded points does not follow the ordering of the originals in $[0, 1]$.

Case 4: Suppose that $L(w) \cup R(w) = \{0, 1\}$, so $L(w) = \{1\}$ and $R(w) = \{0\}$. Let $\{x_i\}$ be the sequence defined in Equation (14) with initial term $x_0 = \frac{1}{2}$. By a similar discussion to Case 1, we have that π is a uniform linear embedding from $[\frac{1}{2}, 1)$ into $\mathbb{R}^{\geq 0}$. To embed the other half of the interval, $[0, \frac{1}{2}]$, let $\{y_i\}$ be the sequence defined in Case 3 with initial term $y_0 = x_1 = r(x_0)$. This is an infinite sequence with limit 0. Now define $\pi'(y_i) = 1 - i$ for all i, $\pi'(y) = 1 - \frac{y - y_0}{y_1 - y_0}$ for $y \in (y_0, y_1)$, and $\pi'(y) = \pi'(r(y)) - 1$ for $y \in (y_i, y_{i+1})$, $i \geq 1$. The map $\pi' : [0, x_1] \to (-\infty, 1]$ is a uniform linear embedding. The maps π and π' agree on $[x_0, x_1]$, which is mapped into $[0, 1]$. Thus, they can be combined to form a uniform embedding from $[0, 1]$ into $\mathbb{R}$.

To show that no uniform embedding exists if $|L(w)| + |R(w)| > 2$, first note that any element $z \in L(w) \cup R(w)$ is associated with an interval $(\ell(z), r(z))$ which is empty, so any uniform embedding must place z at distance at least 1 from all other embedded points. Since $\mathbb{R}$ contains only countably many such points, it follows that, if $L(w) \cap R(w)$ is uncountable, then no uniform embedding exists.

Assume $L(w) \cup R(w)$ is a countable, and therefore nowhere dense subset of $[0,1]$. Let $z \in L(w) \setminus \{1\}$ be an isolated point of $L(w) \cup R(w)$. We can construct an increasing infinite sequence $\{x_i\}_{i=0}^{\infty}$ so that $[x_0, z) \cap (L(w) \cup R(w)) = \emptyset$, and for all i, $x_{i+1} = r(x_i)$. If a uniform linear embedding π exists, then $|\pi(x_{i+1}) - \pi(x_i)| \leq 1$ and $|\pi(x_j) - \pi(x_i)| > 1$ for all $j \geq i+2$. This implies that $\{\pi(x_i)\}_i$ must be either decreasing or increasing. Moreover, since $y > z > r(x)$ for all $y > z > x$, $|\pi(y) - \pi(x_i)| > 1$ for all $y > z$ and $i \geq 0$. Therefore, either $[\pi(x_0), \infty)$ or $(-\infty, \pi(x_0)]$ cannot contain any images from $(z, 1]$.

For every isolated element of $L(w) \cup R(w)$, we can construct such a sequence (increasing for elements of $L(w)$, decreasing for elements of $R(w)$). Moreover, we can choose these sequences such that they fall in disjoint intervals. By the argument above, any sequence "blocks off" a half-line, where no images of the elements of the other sequences can be placed. If there are more than two sequences, this leads to a contradiction. Thus, no uniform linear embedding exists.

$\square$

References

[1] Béla Bollobás, Svante Janson, and Oliver Riordan. The phase transition in inhomogeneous random graphs. *Random Structures and Algorithms*, 31:3–122, 2007.

[2] Béla Bollobás, Svante Janson, and Oliver Riordan. The cut metric, random graphs, and branching processes. *J. Statistical Physics*, 140:289–335, 2010.

[3] Béla Bollobás, Svante Janson, and Oliver Riordan. Monotone graph limits and quasimonotone graphs. *ArXiv e-prints*, January 2011.

[4] Béla Bollobás and Oliver Riordan. Metrics for sparse graphs. In *Surveys in Combinatorics*.

[5] Anthony Bonato, Jeannette Janssen, and Pawel Prałat. Geometric protean graphs. *Internet Mathematics*, 8, 2012.

[6] Chayes J. Lóvasz L. Sós V. T. Borgs, C. and K. Vesztergombi. Convergent sequences of dense graphs i. subgraph frequencies, metric properties and testing. *Adv. Math.*, 219(6):1801–1851.

[7] Chayes J. Lóvasz L. Sós V. T. Borgs, C. and K. Vesztergombi. Convergent sequences of dense graphs ii. multiway cuts and statistical physics. Preprint. Available at http://research.microsoft.com/en-us/um/people/borgs/papers/conright.pdf, 2007.

[8] Christian Borgs, Jennifer Chayes, and László Lovaász. Moments of two-variable functions and the uniqueness of graph limits. *Geometric And Functional Analysis*, 19(6):1597–1619, 2010.

[9] Christian Borgs, Jennifer Chayes, László Lovász, Vera Sós, and Katalin Vesztergombi. Limits of randomly grown graph sequences. *Eur. J. Comb.*, (7):985–999, 2011.

[10] Milan Bradonjic, Arie Hagberg, and Allon G. Percus. The structure of geographical threshold graphs. *Internet Mathematics*, 4:113–139, 2009.

[11] Fan Chung. From quasirandom graphs to graph limits and graphlets. arXiv:1203.2269v6 [math.CO], 2012.

[12] Brent N. Clark, Charles J. Colbourn, and David S. Johnson. Unit disk graphs. *Discrete Math.*, 86(1–3):165–177, 1990.

[13] Corinna Cortes, Daryl Pregibon, and Chris Volinsky. Computational methods for dynamic graphs. *Journal of Computational and Graphical Statistics*, 12.

[14] Jesper Dall and Michael Christensen. Random geometric graphs. *Phys. Rev. E (3)*, 66:1.

[15] Persi Diaconis and Svante Janson. Graph limits and exchangeable random graphs. *Rendiconti di Matematica*, 28:33–61, 2008.

[16] Alan Frieze and Ravi Kannan. Quick approximation to matrices and applications. *Combinatorica*, 19(2):175–220, 1999.

[17] Frédéric Gardi. The Roberts characterization of proper and unit interval graphs. *Discrete Math.*, 307(22):2906–2908, 2007.

[18] Peter D. Hoff, Adrian E. Raftery, and Mark S. Handcock. Latent space approaches to social network analysis. *Journal of the American Statistical Association*, 97(460), 2002.

[19] D. Krioukov, F. Papadopoulos, M. Kitsak, A. Vahdat, and M. Boguñá. Hyperbolic geometry of complex networks. *Phys. Rev. E*, 82:3:036106, 2010.
[20] Jure Leskovec, Jon Kleinberg, and Christos Faloutsos. Graphs over time: Densification laws, shrinking diameters and possible explanations. In *Proc. 11th ACM SIGKDD Intl. Conf. on Knowledge Discovery and Data Mining*, 2005.
[21] László Lovász and Balázs Szegedy. Limits of dense graph sequences. *J. Combin. Theory Ser. B*, (6):933–957, 2006.
[22] László Lovász and Balázs Szegedy. Regularity partitions and the topology of graphons. In *An Irregular Mind: Szemeredi is 70*, pages 415–446. Springer, 2010.
[23] Derek G. Corneil Natasa Pržulj and Igor Jurisica. Modeling interactome: Scale-free or geometric? *Bioinformatics*, 20(18):3508–3515, 2004.
[24] Mathew Penrose. *Random geometric graphs*, volume 5 of *Oxford Studies in Probability*. Oxford University Press, 2003.
[25] Susan Holmes Persi Diaconis and Svante Janson. Interval graph limits. arXiv:1102.2841 [math.CO], to be published in Annals of Combinatorics, 2011.
[26] R.J. Williams and N.D. Martinez. Simple rules yield complex food webs. *Nature*, 393:440–442, 2000.

Department of Mathematics & Statistics, Dalhousie University, Halifax, Nova Scotia, Canada, B3H 3J5